arXiv:1605.06723v1 [math.AP] 22 May 2016

The Boundedness of Bessel-Riesz Operators on Generalized Morrey Spaces

M. Idris[1], H. Gunawan[2], and Eridani[3]

[1,2]Department of Mathematics, Bandung Institute of Technology
Bandung 40132, Indonesia
[3]Department of Mathematics, Airlangga University
Campus C, Mulyorejo, Surabaya 60115, Indonesia
E-mail: [1]mochidris@students.itb.ac.id, [2]hgunawan@math.itb.ac.id, [3]eri.campanato@gmail.com

Abstract. In this paper, we prove the boundedness of Bessel-Riesz operators on generalized Morrey spaces. The proof uses the usual dyadic decomposition, a Hedberg-type inequality for the operators, and the boundedness of Hardy-Littlewood maximal operator. Our results reveal that the norm of the operators is dominated by the norm of the kernels.

Keywords. *Bessel-Riesz operators, Hardy-Littlewood maximal operator, generalized Morrey spaces, boundedness, kernels.*

MSC 2000 Primary 42B20; Secondary 26A33, 42B25, 26D10, 47G10

1 Introduction

We begin with the definition of Bessel-Riesz operators. For $0 < \gamma$ and $0 < \alpha < n$, we define

$$I_{\alpha,\gamma} f(x) := \int_{\mathbb{R}^n} K_{\alpha,\gamma}(x-y) f(y)\, dy,$$

for every $f \in L^p_{loc}(\mathbb{R}^n)$, $p \geq 1$, where $K_{\alpha,,\gamma}(x) := \frac{|x|^{\alpha-n}}{(1+|x|)^{\gamma}}$. Here, $I_{\alpha,\gamma}$ is called *Bessel-Riesz operator* and $K_{\alpha,\gamma}$ is called *Bessel-Riesz kernel*. The name of the kernel resembles the product of Bessel kernel and Riesz kernel [13]. While the Riesz kernel captures the local behaviour, the Bessel kernel take cares the global behaviour of the function. The Bessel-Riesz kernel is used in studying the behaviour of the solution of a Schrödinger type equation [8].

For $\gamma = 0$, we have $I_{\alpha,0} = I_\alpha$, known as *fractional integral operators* or *Riesz potentials* [12]. Around 1930, Hardy and Littlewood [5, 6] and Sobolev [11] have proved the boundedness of I_α on Lebesgue spaces via the inequality

$$\|I_\alpha f\|_{L^q} \leq C \, \|f\|_{L^p},$$

for every $f \in L^p(\mathbb{R}^n)$, where $1 < p < \frac{\alpha}{n}$, and $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$. Here C denotes a constant which may depend on α, p, q, and n, but not on f.

For $1 \leq p \leq q$, the *(classical) Morrey space* $L^{p,q}(\mathbb{R}^n)$ is defined by

$$L^{p,q}(\mathbb{R}^n) := \{f \in L^p_{loc}(\mathbb{R}^n) : \|f\|_{L^{p,q}} < \infty\},$$

where $\|f\|_{L^{p,q}} := \sup_{r>0,a\in\mathbb{R}^n} r^{n(1/q-1/p)} \left(\int_{|x-a|<r} |f(x)|^p dx\right)^{1/p}$. For these spaces, we have an inclusion property which is presented by the following theorem.

Theorem 1.1 *For $1 \le p \le q$, we have $L^q(\mathbb{R}^n) = L^{q,q}(\mathbb{R}^n) \subseteq L^{p,q}(\mathbb{R}^n) \subseteq L^{1,q}(\mathbb{R}^n)$.*

On Morrey spaces, Spanne [10] has shown that I_α is bounded form $L^{p_1,q_1}(\mathbb{R}^n)$ to $L^{p_2,q_2}(\mathbb{R}^n)$ for $1 < p_1 < q_1 < \frac{n}{\alpha}$, $\frac{1}{p_2} = \frac{1}{p_1} - \frac{\alpha}{n}$, and $\frac{1}{q_2} = \frac{1}{q_1} - \frac{\alpha}{n}$. Furthermore, Adams [1] and Chiarenza and Frasca [2] reproved it and obtained a stronger result which is presented by the following theorem.

Theorem 1.2 *[Adams, Chiarenza-Frasca] If $0 < \alpha < n$, then we have*

$$\|I_\alpha f\|_{L^{p_2,q_2}} \le C \|f\|_{L^{p_1,q_1}},$$

for every $f \in L^{p_1,q_1}(\mathbb{R}^n)$ where $1 < p_1 < q_1 < \frac{n}{\alpha}$, $\frac{1}{p_2} = \frac{1}{p_1}\left(1 - \frac{\alpha q_1}{n}\right)$, and $\frac{1}{q_2} = \frac{1}{q_1} - \frac{\alpha}{n}$.

For $\phi : \mathbb{R}^+ \to \mathbb{R}^+$ and $1 \le p < \infty$, we define the *generalized Morrey space*

$$L^{p,\phi}(\mathbb{R}^n) := \{f \in L^p_{loc}(\mathbb{R}^n) : \|f\|_{L^{p,\phi}} < \infty\},$$

where $\|f\|_{L^{p,\phi}} := \sup_{r>0,a\in\mathbb{R}^n} \frac{1}{\phi(r)} \left(\frac{1}{r^n} \int_{|x-a|<r} |f(x)|^p \, dx\right)^{1/p}$. Here we assume that ϕ is almost decreasing and $t^{n/p}\phi(t)$ is almost decreasing, so that ϕ satisfies the *doubling condition*, that is, there exists a constant C such that $\frac{1}{C} \le \frac{\phi(r)}{\phi(v)} \le C$ whenever $\frac{1}{2} \le \frac{r}{v} \le 2$.

In 1994, Nakai [9] obtained the boundedness of I_α from $L^{p_1,\phi}(\mathbb{R}^n)$ to $L^{p_2,\psi}(\mathbb{R}^n)$ where $1 < p_1 < \frac{n}{\alpha}$, $\frac{1}{p_2} = \frac{1}{p_1} - \frac{\alpha}{n}$ and $\int_r^\infty v^{\alpha-1}\phi(v)dv \le Cr^\alpha\phi(r) \le C\psi(r)$ for every $r > 0$. Nakai's result may be viewed as an extension of Spanne's. Later on, in 2009, Gunawan and Eridani [3] extended Adams-Chiarenza-Frasca's result.

Theorem 1.3 *[Gunawan-Eridani] If $\int_r^\infty \frac{\phi(v)}{v} dv \le C\phi(r)$, and $\phi(r) \le Cr^\beta$ for every $r > 0$, $-\frac{n}{p_1} \le \beta < -\alpha$, $1 < p_1 < \frac{n}{\alpha}$, $0 < \alpha < n$, then we have*

$$\|I_\alpha f\|_{L^{p_2,\psi}} \le C \|f\|_{L^{p_1,\phi}}$$

for every $f \in L^{p_1,\phi}(\mathbb{R}^n)$ where $p_2 = \frac{\beta p_1}{\alpha+\beta}$ and $\psi(r) = \phi(r)^{p_1/p_2}$, $r > 0$.

The proof of the boundedness of I_α on Lebesgue spaces, Morrey spaces, or generalized Morrey spaces, usually involves *Hardy-Littlewood maximal operator*, which is defined by

$$Mf(x) := \sup_{B\ni x} \frac{1}{|B|} \int_B |f(y)| \, dy,$$

for every $f \in L^p_{loc}(\mathbb{R}^n)$ where $|B|$ denotes the Lebesgue measure of the ball $B = B(a,r)$ (centered at $a \in \mathbb{R}^n$ with radius $r > 0$). It is well known that the operator M is bounded on $L^p(\mathbb{R}^n)$ for $1 < p \le \infty$ [12, 13] and also on Morrey spaces $L^{p,q}$ for $1 < p \le q \le \infty$ [2].

Next, we know that $I_{\alpha,\gamma}$ is guaranteed to be bounded on generalized Morrey spaces because $K_{\alpha,\gamma}(x) \le K_\alpha(x)$ for every $x \in \mathbb{R}^n$. The boundedness of $I_{\alpha,\gamma}$ on Lebesgue spaces can also be proved by using Young's inequality, as shown in [7].

Theorem 1.4 *[7] For $0 < \gamma$ and $0 < \alpha < n$, we have $K_{\alpha,\gamma} \in L^t(\mathbb{R}^n)$ whenever $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$. Accordingly, we have*

$$\|I_{\alpha,\gamma} f\|_{L^q} \leq \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^p}$$

for every $f \in L^p(\mathbb{R}^n)$ where $1 \leq p < t'$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, $\frac{1}{q} + 1 = \frac{1}{p} + \frac{1}{t}$.

Using the boundedness of Hardy-Littlewood maximal operator, we also know that $I_{\alpha,\gamma}$ is bounded on Morrey spaces.

Theorem 1.5 *[7] For $0 < \gamma$ and $0 < \alpha < n$, we have*

$$\|I_{\alpha,\gamma} f\|_{L^{p_2,q_2}} \leq C \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,q_1}}$$

for every $f \in L^{p_1,q_1}(\mathbb{R}^n)$ where $1 < p_1 < q_1 < t'$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, $\frac{1}{p_2} + 1 = \frac{1}{p_1} + \frac{1}{t}$, and $\frac{1}{q_2} + 1 = \frac{1}{q_1} + \frac{1}{t}$.

In 1999, Kurata *et al.* [8] proved the boundedness of $W \cdot I_{\alpha,\gamma}$ on generalized Morrey spaces where W is a multiplication operator. A similar result to Kurata's can be found in [3]. In the next section, we shall reprove the boundedness of $I_{\alpha,\gamma}$ on generalized Morrey spaces using a Hedberg-type inequality and the boundedness of Hardy-Littlewood maximal operator on these spaces.

Theorem 1.6 *(Nakai) For $1 < p \leq \infty$, we have*

$$\|Mf\|_{L^{p,\psi}} \leq C \|f\|_{L^{p,\phi}},$$

for every $f \in L^{p,\phi}(\mathbb{R}^n)$.

Our results show that the norm of Bessel-Riesz operators is dominated by the norm of their kernels on (generalized) Morrey spaces.

2 Inequalities For $I_{\alpha,\gamma}$ On Generalized Morrey Spaces

For $0 < \alpha < n$ and $\gamma > 0$, one may observe that the kernel $K_{\alpha,\gamma}$ belongs to Lebesgue spaces $L^t(\mathbb{R}^n)$ whenever $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, where

$$\sum_{k\in\mathbb{Z}} \frac{(2^k R)^{(\alpha-n)t+n}}{(1+2^k R)^{\gamma t}} \sim \|K_{\alpha,\gamma}\|_{L^t}^t$$

(see [7]). With this in mind, we obtain the boundedness of $I_{\alpha,\gamma}$ on generalized Morrey spaces as in the following theorem.

Theorem 2.1 *Let $0 < \gamma$ and $0 < \alpha < n$. If $\phi(r) \leq Cr^\beta$ for every $r > 0$, $-\frac{\alpha t'}{p_1} \leq \beta < -\alpha$, $1 < p_1 < t'$, and $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, then we have*

$$\|I_{\alpha,\gamma} f\|_{L^{p_2,\psi}} \leq C \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,\phi}},$$

for every $f \in L^{p_1,\phi}(\mathbb{R}^n)$ where $p_2 = \frac{\beta p_1}{\alpha+\beta}$, and $\psi(r) = \phi(r)^{p_1/p_2}$.

Proof. Let $0 < \gamma$ and $0 < \alpha < n$. Suppose that $\phi(r) \leq Cr^{\beta}$ for every $r > 0$, $-\frac{\alpha t'}{p_1} \leq \beta < -\alpha$, $1 < p_1 < t'$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$. Take $f \in L^{p_1,\phi}(\mathbb{R}^n)$ and write

$$I_{\alpha,\gamma} f(x) := I_1(x) + I_2(x),$$

for every $x \in \mathbb{R}^n$ where $I_1(x) := \int_{|x-y|<R} \frac{|x-y|^{\alpha-n} f(y)}{(1+|x-y|)^{\gamma}} dy$ and $I_2(x) := \int_{|x-y|\geq R} \frac{|x-y|^{\alpha-n} f(y)}{(1+|x-y|)^{\gamma}} dy$, $R > 0$.

Using dyadic decomposition, we have the following estimate for I_1:

$$\begin{aligned} |I_1(x)| &\leq \sum_{k=-\infty}^{-1} \int_{2^k R \leq |x-y| < 2^{k+1}R} \frac{|x-y|^{\alpha-n} |f(y)|}{(1+|x-y|)^{\gamma}} dy \\ &\leq C_1 \sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{\alpha-n}}{\left(1+2^k R\right)^{\gamma}} \int_{2^k R \leq |x-y| < 2^{k+1}R} |f(y)| \, dy \\ &\leq C_2 \, Mf(x) \sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{\alpha-n+n/t} \left(2^k R\right)^{n/t'}}{\left(1+2^k R\right)^{\gamma}}. \end{aligned}$$

We then use Hölder's inequality to get

$$|I_1(x)| \leq C_2 Mf(x) \left(\sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{(\alpha-n)t+n}}{\left(1+2^k R\right)^{\gamma t}} \right)^{1/t} \left(\sum_{k=-\infty}^{-1} \left(2^k R\right)^{n} \right)^{1/t'}.$$

Because we have

$$\left(\sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{(\alpha-n)t+n}}{\left(1+2^k R\right)^{\gamma t}} \right)^{1/t} \leq \left(\sum_{k\in\mathbb{Z}} \frac{\left(2^k R\right)^{(\alpha-n)t+n}}{\left(1+2^k R\right)^{\gamma t}} \right)^{1/t} \sim \|K_{\alpha,\gamma}\|_{L^t},$$

we obtain $|I_1(x)| \leq C_3 \|K_{\alpha,\gamma}\|_{L^t} Mf(x) R^{n/t'}$.

To estimate I_2, we use Hölder's inequality again:

$$\begin{aligned} |I_2(x)| &\leq C_4 \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha-n}}{\left(1+2^k R\right)^{\gamma}} \int_{2^k R \leq |x-y| < 2^{k+1}R} |f(y)| \, dy \\ &\leq C_4 \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha-n}}{\left(1+2^k R\right)^{\gamma}} \left(2^k R\right)^{n/p_1'} \left(\int_{2^k R \leq |x-y| < 2^{k+1}R} |f(y)|^{p_1} \, dy \right)^{1/p_1}. \end{aligned}$$

It follows that

$$\begin{aligned} |I_2(x)| &\leq C_5 \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha-n+n/t}}{\left(1+2^k R\right)^{\gamma}} \phi\left(2^k R\right) \left(2^k R\right)^{n/t'} \\ &\leq C_6 \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha-n+n/t}}{\left(1+2^k R\right)^{\gamma}} \left(2^k R\right)^{\beta+n/t'}. \end{aligned}$$

Another use of Hölder's inequality gives

$$|I_2(x)| \leq C_6 \|f\|_{L^{p_1,\phi}} \left(\sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{(\alpha-n)t+n}}{\left(1+2^k R\right)^{\gamma t}} \right)^{1/t} \left(\sum_{k=0}^{\infty} \left(2^k R\right)^{\beta t'+n} \right)^{1/t'}.$$

Because $\beta t' + n < 0$ and $\sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{(\alpha-n)t+n}}{\left(1+2^k R\right)^{\gamma t}} \lesssim \|K_{\alpha,\gamma}\|_{L^t}^t$, we obtain

$$|I_2(x)| \leq C_7 \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,\phi}} R^{\beta} R^{n/t'}.$$

Summing the two estimates, we obtain

$$|I_{\alpha,\gamma}f(x)| \leq C_8 \|K_{\alpha,\gamma}\|_{L^t} \left(Mf(x) R^{n/t'} + \|f\|_{L^{p_1,\phi}} R^{n/t'+\beta} \right),$$

for every $x \in \mathbb{R}^n$. Now, for each $x \in \mathbb{R}^n$, choose $R > 0$ such that $R^{-\beta} = \frac{\|f\|_{L^{p_1,\phi}}}{Mf(x)}$. Hence we get a Hedberg-type inequality for $I_{\alpha,\gamma}f$, namely

$$|I_{\alpha,\gamma}f(x)| \leq C_9 \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,\phi}}^{-\alpha/\beta} Mf(x)^{1+\alpha/\beta}.$$

Now put $p_2 := \frac{\beta p_1}{\alpha+\beta}$. For arbitrary $a \in \mathbb{R}^n$ and $r > 0$, we have

$$\left(\int_{|x-a|<r} |I_{\alpha,\gamma}f(x)|^{p_2} dx \right)^{1/p_2} \leq C_9 \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \left(\int_{|x|<r} |Mf(x)|^{p_1} dx \right)^{1/p_2}.$$

We divide both sides by $\phi(r)^{p_1/p_2} r^{n/p_2}$ to get

$$\frac{\left(\int_{|x|<r} |I_{\alpha,\gamma}f(x)|^{p_2} dx \right)^{1/p_2}}{\psi(r) r^{n/p_2}} \leq C_9 \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \frac{\left(\int_{|x|<r} |Mf(x)|^{p_1} dx \right)^{(1/p_2)}}{\phi(r)^{p_1/p_2} r^{n/p_2}},$$

where $\psi(r) := \phi(r)^{p_1/p_2}$. Taking the supremum over $a \in \mathbb{R}^n$ and $r > 0$, we obtain

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C_{10} \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \|Mf\|_{L^{p_1,\phi}}^{p_1/p_2}.$$

By the boundedness of the maximal operator on generalized Morrey spaces (Nakai's Theorem), the desired result follows: $\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C \|K_{\alpha,\gamma}\|_{L^t} \|f\|_{L^{p_1,\phi}}$. ■

We note that from the inclusion property of Morrey spaces, we have $\|I_{\alpha,\gamma}f\|_{L^{s,t}} \leq \|I_{\alpha,\gamma}f\|_{L^{t,t}} = \|I_{\alpha,\gamma}f\|_{L^t}$ where $1 \leq s \leq t$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$. Now we wish to find a better upper bound for the norm of $I_{\alpha,\gamma}$. For this purpose, we shall use the fact that the kernel $K_{\alpha,\gamma}$ belongs to Morrey spaces.

Theorem 2.2 *Let $0 < \gamma$ and $0 < \alpha < n$. If $\phi(r) \leq Cr^\beta$ for every $r > 0$, $-\frac{\alpha t'}{p_1} \leq \beta < -\alpha$, $1 < p_1 < t'$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$, then we have*

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}},$$

for every $f \in L^{p_1,\phi}(\mathbb{R}^n)$ where $1 \leq s \leq t$, $p_2 = \frac{\beta p_1}{\alpha+\beta}$, and $\psi(v) = \phi(v)^{p_1/p_2}$.

Proof. Let $0 < \gamma$ and $0 < \alpha < n$. Suppose that $\phi(r) \leq Cr^\beta$ for every $r > 0$, $-\frac{\alpha t'}{p_1} \leq \beta < -\alpha$, $1 < p_1 < t'$, $\frac{n}{n+\gamma-\alpha} < t < \frac{n}{n-\alpha}$. As in the proof of Theorem 2.1, we have $I_{\alpha,\gamma}f(x) := I_1(x) + I_2(x)$ for every $x \in \mathbb{R}^n$. Now, we estimate I_1 using dyadic decomposition as follow:

$$\begin{aligned}
|I_1(x)| &\leq \sum_{k=-\infty}^{-1} \int_{2^k R \leq |x-y| < 2^{k+1}R} \frac{|x-y|^{\alpha-n} |f(y)|}{(1+|x-y|)^\gamma} dy \\
&\leq C_1 \sum_{k=-\infty}^{-1} \frac{(2^k R)^{\alpha-n}}{(1+2^k R)^\gamma} \int_{2^k R \leq |x-y| < 2^{k+1}R} |f(y)| dy \\
&= C_2 Mf(x) \sum_{k=-\infty}^{-1} \frac{(2^k R)^{\alpha-n+n/s} (2^k R)^{n/s'}}{(1+2^k R)^\gamma},
\end{aligned}$$

where $1 \leq s \leq t$. By Hölder's inequality,

$$|I_1(x)| \leq C_2 Mf(x) \left(\sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{(\alpha-n)s+n}}{\left(1+2^k R\right)^{\gamma s}} \right)^{1/s} \left(\sum_{k=-\infty}^{-1} \left(2^k R\right)^n \right)^{1/s'}.$$

We also have $\sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{(\alpha-n)s+n}}{\left(1+2^k R\right)^{\gamma s}} \lesssim \int_{0<|x|<R} K_{\alpha,\gamma}^s(x)\,dx$, so that

$$|I_1(x)| \leq C_3 Mf(x) \left(\int_{0<|x|<R} K_{\alpha,\gamma}^s(x)\,dx \right)^{\frac{1}{s}} R^{n/s'} \leq C_3 \|K_{\alpha,\gamma}\|_{L^{s,t}} Mf(x) R^{n/t'}.$$

Next, we estimate I_2 by using Hölder's inequality. As in the proof of Theorem 2.1, we obtain

$$|I_2(x)| \leq C_4 \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha-n}}{\left(1+2^k R\right)^{\gamma}} \left(2^k R\right)^{n/p_1'} \left(\int_{2^k R \leq |x-y| < 2^{k+1} R} |f(y)|^{p_1}\,dy \right)^{1/p_1}.$$

It thus follows that

$$\begin{aligned} |I_2(x)| &\leq C_5 \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha} \phi(2^k R)}{\left(1+2^k R\right)^{\gamma}} \frac{\left(\int_{2^k R \leq |x-y| < 2^{k+1} R} dy \right)^{1/s}}{\left(2^k R\right)^{n/s}} \\ &\leq C_6 \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \phi(2^k R) \left(2^k R\right)^{n/t'} \frac{\left(\int_{2^k R \leq |x-y| < 2^{k+1} R} K_{\alpha,\gamma}^s(x-y) dy \right)^{1/s}}{\left(2^k R\right)^{n/s-n/t}} \end{aligned}$$

Because $\phi(r) \leq C r^{\beta}$ and $\frac{\left(\int_{2^k R \leq |x-y| < 2^{k+1} R} K_{\alpha,\gamma}^s(x-y) dy \right)^{1/s}}{\left(2^k R\right)^{n/s-n/t}} \lesssim \|K_{\alpha,\gamma}\|_{L^{s,t}}$ for every $k = 0, 1, 2, \ldots$, we get

$$\begin{aligned} |I_2(x)| &\leq C_7 \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \left(2^k R\right)^{\beta+n/t'} \\ &\leq C_8 \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}} R^{\beta} R^{n/t'}. \end{aligned}$$

From the two estimates, we obtain

$$|I_{\alpha,\gamma} f(x)| \leq C_9 \|K_{\alpha,\gamma}\|_{L^{s,t}} \left(Mf(x) R^{n/t'} + \|f\|_{L^{p_1,\phi}} R^{n/t'+\beta} \right),$$

for every $x \in \mathbb{R}^n$. Now, for each $x \in \mathbb{R}^n$, choose $R > 0$ such that $R^{-\beta} = \frac{\|f\|_{L^{p_1,\phi}}}{Mf(x)}$. Hence we get

$$|I_{\alpha,\gamma} f(x)| \leq C_9 \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}}^{-\alpha/\beta} Mf(x)^{1+\alpha/\beta}.$$

Put $p_2 := \frac{\beta p_1}{\alpha+\beta}$. For arbitrary $a \in \mathbb{R}^n$ and $r > 0$, we have

$$\left(\int_{|x-a|<r} |I_{\alpha,\gamma} f(x)|^{p_2}\,dx \right)^{1/p_2} \leq C_9 \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \left(\int_{|x-a|<r} |Mf(x)|^{p_1}\,dx \right)^{1/p_2}.$$

Divide both sides by $\phi(r)^{p_1/p_2} r^{n/p_2}$ and take the supremum over $a \in \mathbb{R}^n$ and $r > 0$ to get

$$\|I_{\alpha,\gamma} f\|_{L^{p_2,\psi}} \leq C_{10} \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \|Mf\|_{L^{p_1,\phi}}^{p_1/p_2},$$

where $\psi(r) := \phi(r)^{p_1/p_2}$. With the boundedness of the maximal operator on generalized Morrey spaces (Nakai's Theorem), we obtain the desired result: $\|I_{\alpha,\gamma} f\|_{L^{p_2,\psi}} \leq C_{p_1,\phi} \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}}$. ■

Note that by Theorem 2.2 and the inclusion of Morrey spaces, we recover Theorem 2.1:

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C\,\|K_{\alpha,\gamma}\|_{L^{s,t}}\,\|f\|_{L^{p_1,\phi}} \leq C\,\|K_{\alpha,\gamma}\|_{L^t}\,\|f\|_{L^{p_1,\phi}}.$$

We wish to obtain a better estimate. The following lemma presents that the Bessel-Riesz kernels belong to generalized Morrey space $L^{s,\sigma}(\mathbb{R}^n)$ for some $s \geq 1$ and a suitable function σ.

Lemma 2.3 *Suppose that $0 < \gamma$ and $0 < \alpha < n$. If $\sigma : \mathbb{R}^+ \to \mathbb{R}^+$ satisfies $\int_{0<r\leq R} r^{(\alpha-n)s+n-1}dr \leq C\sigma^s(R)R^n$ for every $R > 0$, then $K_{\alpha,\gamma} \in L^{s,\sigma}(\mathbb{R}^n)$.*

Proof. Suppose that the hypothesis holds. We observe that

$$\int_{|x-0|\leq R} K^s_{\alpha,\gamma}(x)dx \;=\; \int_{|x|\leq R} \frac{|x|^{(\alpha-n)s}}{(1+|x|)^{\gamma s}}dx \leq C\int_{0<r\leq R} r^{(\alpha-n)s+n-1}dr \leq C\sigma^s(R)R^n.$$

We divide both sides of the inequality by $\sigma^s(R)R^n$ and take s^{th}-root to obtain $\frac{\left(\int_{|x-0|\leq R} K^s_{\alpha,\gamma}(x)dx\right)^{1/s}}{\sigma(R)R^{n/s}} \leq C^{1/s}$. Now, taking the supremum over $R > 0$, we have $\sup\limits_{R>0} \frac{\left(\int_{|x-0|\leq R} K^s_{\alpha,\gamma}(x)dx\right)^{1/s}}{\sigma(R)R^{n/s}} < \infty$. Hence $K_{\alpha,\gamma} \in L^{s,\sigma}(\mathbb{R}^n)$. ■

By the hypothesis of Lemma 2.3 we also obtain $\frac{\left(\int_{2^kR<|x|\leq 2^{k+1}R} K^s_{\rho,\gamma}(x)dx\right)^{1/s}}{\sigma(2^kR)(2^kR)^{n/s}} \lesssim \|K_{\alpha,\gamma}\|_{L^{s,\sigma}}$ for every integer k and $R > 0$. Moreover, $\frac{\left(\sum_{k=-1}^{-\infty} K^s_{\alpha,\gamma}(2^kR)(2^kR)^n\right)^{1/s}}{\sigma(R)(R)^{n/s}} \lesssim \|K_{\alpha,\gamma}\|_{L^{s,\sigma}}$ holds for every $R > 0$. One may observe that $1 \leq s \leq \frac{n\ln R_1}{-\ln\sigma(R_1)}$ for every $R_1 > 1$. For $\sigma(R) = R^{-n/t}$, this inequality reduces to $1 \leq s \leq t$.

We shall now use the lemma to prove the following theorem.

Theorem 2.4 *Suppose that $\sigma : \mathbb{R}^+ \to \mathbb{R}^+$ satisfies the doubling condition and $\sigma(r) \leq Cr^{-\alpha}$ for every $r > 0$, so that $K_{\alpha,\gamma} \in L^{s,\sigma}(\mathbb{R}^n)$ for $1 \leq s < \frac{n}{n-\alpha}$, where $0 < \alpha < n$ and $\gamma > 0$. If $\phi(r) \leq Cr^{\beta}$ for every $r > 0$, where $-\frac{n}{p_1} < \beta < -\alpha$, then we have*

$$\|I_{\alpha,\gamma}f\|_{L^{p_2,\psi}} \leq C_{p_1,\phi}\,\|K_{\alpha,\gamma}\|_{L^{s,\sigma}}\,\|f\|_{L^{p_1,\phi}}$$

for every $f \in L^{p_1,\phi}(\mathbb{R}^n)$, where $1 < p_1 < \frac{n}{\alpha}$, $p_2 = \frac{\beta p_1}{\beta+n-\alpha}$ and $\psi(r) = \phi(r)^{p_1/p_2}$.

Proof. Let $0 < \gamma$ and $0 < \alpha < n$. Suppose that $\sigma : \mathbb{R}^+ \to \mathbb{R}^+$ satisfies the doubling condition and $\sigma(r) \leq Cr^{-\alpha}$ for every $r > 0$, such that $K_{\alpha,\gamma} \in L^{s,\sigma}(\mathbb{R}^n)$ for $1 \leq s < \frac{n}{n-\alpha}$. Suppose also that $\phi(r) \leq Cr^{\beta}$ for every $r > 0$, where $-\frac{n}{p_1} < \beta < -\alpha$, $1 < p_1 < \frac{n}{\alpha}$. As in the proof of Theorem 2.1, we write $I_{\alpha,\gamma}f(x) := I_1(x) + I_2(x)$ for every $x \in \mathbb{R}^n$. As usual, we estimate I_1 by using dyadic decomposition:

$$\begin{aligned}
|I_1(x)| &\leq \sum_{k=-\infty}^{-1} \int_{2^kR\leq|x-y|<2^{k+1}R} \frac{|x-y|^{\alpha-n}|f(y)|}{(1+|x-y|)^{\gamma}}dy \\
&\leq C_1 \sum_{k=-\infty}^{-1} \frac{(2^kR)^{\alpha-n}}{(1+2^kR)^{\gamma}} \int_{2^kR\leq|x-y|<2^{k+1}R} |f(y)|\,dy \\
&= C_2 Mf(x) \sum_{k=-\infty}^{-1} \frac{(2^kR)^{\alpha-n+n/s}(2^kR)^{n/s'}}{(1+2^kR)^{\gamma}}
\end{aligned}$$

By using Hölder inequality, we obtain

$$|I_1(x)| \leq C_2 Mf(x) \left(\sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{(\alpha-n)s+n}}{\left(1+2^k R\right)^{\gamma s}} \right)^{1/s} \left(\sum_{k=-\infty}^{-1} \left(2^k R\right)^n \right)^{1/s'}.$$

But $\sum_{k=-\infty}^{-1} \frac{\left(2^k R\right)^{(\alpha-n)s+n}}{\left(1+2^k R\right)^{\gamma s}} \lesssim \int_{0<|x|<R} K_{\alpha,\gamma}^s(x)\,dx$, and so we get

$$\begin{aligned} |I_1(x)| &\leq C_2 Mf(x) \left(\int_{0<|x|<R} K_{\alpha,\gamma}^s(x)\,dx \right)^{\frac{1}{s}} R^{n/s'} \\ &\leq C_2 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} Mf(x)\,\sigma(R)\,R^n \\ &\leq C_3 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} Mf(x)\,R^{n-\alpha}. \end{aligned}$$

Next, we estimate I_2 as follows:

$$\begin{aligned} |I_2(x)| &\leq C_4 \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha-n}}{\left(1+2^k R\right)^{\gamma}} \int_{2^k R\leq|x-y|<2^{k+1}R} |f(y)|\,dy \\ &\leq C_4 \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha-n}}{\left(1+2^k R\right)^{\gamma}} \left(2^k R\right)^{n/p_1'} \left(\int_{2^k R\leq|x-y|<2^{k+1}R} |f(y)|^{p_1}\,dy \right)^{1/p_1} \\ &\leq C_5 \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \frac{\left(2^k R\right)^{\alpha} \phi\left(2^k R\right) \left(2^k R\right)^n}{\left(1+2^k R\right)^{\gamma}} \frac{\left(\int_{2^k R\leq|x-y|<2^{k+1}R} dy\right)^{1/s}}{\left(2^k R\right)^{n/s}} \\ &\leq C_6 \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \left(2^k R\right)^{n-\alpha+\beta} \frac{\left(\int_{2^k R\leq|x-y|<2^{k+1}R} K_{\alpha,\gamma}^s(x-y)\,dy\right)^{1/s}}{\sigma\left(2^k R\right)\left(2^k R\right)^{n/s}} \end{aligned}$$

Because $\frac{\int_{2^k R\leq|x-y|<2^{k+1}R} K_{\alpha,\gamma}^s(x-y)dy}{\sigma(2^k R)(2^k R)^n} \lesssim \|K_{\alpha,\gamma}\|_{L^{s,\sigma}}^s$ for every $k = 0, 1, 2, \ldots$, we obtain

$$\begin{aligned} |I_2(x)| &\leq C_6 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}} \sum_{k=0}^{\infty} \left(2^k R\right)^{n-\alpha+\beta} \\ &\leq C_7 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}} R^{n-\alpha+\beta}. \end{aligned}$$

It follows from the above estimates for I_1 and I_2 that

$$|I_{\alpha,\gamma} f(x)| \leq C_8 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \left(Mf(x)\,R^{n-\alpha} + \|f\|_{L^{p_1,\phi}} R^{n-\alpha+\beta} \right)$$

for every $x \in \mathbb{R}^n$. Now, for each $x \in \mathbb{R}^n$, choose $R > 0$ such that $R^{-\beta} = \frac{\|f\|_{L^{p_1,\phi}}}{Mf(x)}$, whence

$$|I_{\alpha,\gamma} f(x)| \leq C_9 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}}^{(\alpha-n)/\beta} Mf(x)^{1+(n-\alpha)/\beta}.$$

Put $p_2 := \frac{\beta p_1}{\beta+n-\alpha}$. For arbitrary $a \in \mathbb{R}^n$ and $r > 0$, we have

$$\left(\int_{|x-a|<r} |I_{\alpha,\gamma} f(x)|^{p_2}\,dx \right)^{1/p_2} \leq C_9 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \left(\int_{|x-a|<r} |Mf(x)|^{p_1}\,dx \right)^{(1/p_2)}.$$

Divide the both sides by $\phi(r)^{p_1/p_2} r^{n/p_2}$ to get

$$\begin{aligned} \frac{\left(\int_{|x|<r} |I_{\alpha,\gamma} f(x)|^{p_2}\,dx\right)^{1/p_2}}{\psi(r)\,r^{n/p_2}} &= \frac{\left(\int_{|x|<r} |I_{\alpha,\gamma} f(x)|^{p_2}\,dx\right)^{1/p_2}}{\phi(r)^{p_1/p_2}\,r^{n/p_2}} \\ &\leq C_9 \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \frac{\left(\int_{|x|<r} |Mf(x)|^{p_1}\,dx\right)^{(1/p_2)}}{\phi(r)^{p_1/p_2}\,r^{n/p_2}}, \end{aligned}$$

where $\psi(r) := \phi(r)^{p_1/p_2}$. Finally, take the supremum over $a \in \mathbb{R}^n$ and $r > 0$ to obtain

$$\|I_{\alpha,\gamma} f\|_{L^{p_2,\psi}} \leq C_{10} \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}}^{1-p_1/p_2} \|Mf\|_{L^{p_1,\phi}}^{p_1/p_2}.$$

Because the maximal operator is bounded on generalized Morrey spaces (Nakai's Theorem), we conclude that $\|I_{\alpha,\gamma} f\|_{L^{p_2,\psi}} \leq C_{p_1,\phi} \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}}$. ■

3 Concluding Remarks

The results presented in this paper, namely Theorems 2.1, 2.2, and 2.4, extend the results on the boundedness of Bessel-Riesz operators on Morrey spaces [7]. Similar to Gunawan-Eridani's result for I_α, Theorems 2.1, 2.2, and 2.4 ensures that $I_{\alpha,\gamma} : L^{p_1,\phi}(\mathbb{R}^n) \to L^{p_2,\phi^{p_1/p_2}}(\mathbb{R}^n)$. Notice that if we have $\sigma : \mathbb{R}^+ \to \mathbb{R}^+$ such that for $t \in (\frac{n}{n+\gamma-\alpha}, \frac{n}{n-\alpha})$, then $R^{-n/t} < \sigma(R)$ holds for every $R > 0$, then Theorem 2.4 gives a better estimate than Theorem 2.2. Now, if we define $\sigma(R) := \left(1 + R^{n/t_1}\right) R^{-n/t}$ for some $t_1 > t$, then $\|K_{\alpha,\gamma}\|_{L^{s,\sigma}} < \|K_{\alpha,\gamma}\|_{L^{s,t}}$. By Theorem 2.2 and the inclusion property of Morrey spaces, we obtain

$$\begin{aligned} \|I_{\alpha,\gamma} f\|_{L^{p_2,\psi}} &\leq C \|K_{\alpha,\gamma}\|_{L^{s,\sigma}} \|f\|_{L^{p_1,\phi}} \\ &< C \|K_{\alpha,\gamma}\|_{L^{s,t}} \|f\|_{L^{p_1,\phi}} \\ &\leq C \|K_{\alpha,\gamma}\|_{L^{t}} \|f\|_{L^{p_1,\phi}}. \end{aligned}$$

We can therefore say that Theorem 2.4 gives the best estimate among the three. Furthermore, we have also shown that, in each theorem, the norm of Bessel-Riesz operators on generalized Morrey spaces is dominated by that of Bessel-Riesz kernels.

Acknowledgements. The first and second authors are supported by ITB Research & Innovation Program 2015.

References

[1] D.R. Adams, "A note on Riesz potentials", *Duke Math. J.* **42** (1975), 765-778.

[2] F. Chiarenza and M. Frasca, "Morrey spaces and Hardy-Littlewood maximal function", *Rend. Mat.* **7** (1987), 273-279.

[3] H. Gunawan and Eridani, "Fractional integrals and generalized Olsen inequalities", *Kyungpook Math. J.* **1** (2009) 49.

[4] L. Grafakos, *Classical Fourier Analysis*, Graduate Texts in Matemathics, Vol. 249, Springer, New York, 2008.

[5] G.H. Hardy and J.E. Littlewood, "Some properties of fractional integrals. I", *Math. Zeit.* **27** (1927), 565-606.

[6] G.H. Hardy and J.E. Littlewood, "Some properties of fractional integrals. II", *Math. Zeit.* **34** (1932), 403-439.

[7] M. Idris, H. Gunawan, J. Lindiarni, and Eridani, "The boundedness of Bessel-Riesz operators on Morrey spaces", submitted to the Proceedings of ISCPMS 2015.

[8] K. Kurata, S. Nishigaki and S. Sugano, "Boundedness of integral operator on generalized Morrey spaces and its aplication to Schrödinger operator", *Proc. Amer. Math. Soc.* **128** (1999), 587-602.

[9] E. Nakai, "Hardy-Littlewood maximal operator, singular integral operator and Riesz potentials on generalized Morrey spaces", *Math. Nachr.* **166** (1994), 95-103.

[10] J. Peetre, "On the theory of $\mathcal{L}_{p,\lambda}$ spaces", *J. Funct. Anal.* **4** (1969), 71-87.

[11] S.L. Sobolev, "On a theorem in functional analysis" (Russian), *Math. Sob.* **46** (1938), 471-497. [English translation in *Amer. Math. Soc. Transl. ser.2.* **34** (1963), 39-68].

[12] E. M. Stein, *Singular Integral and Differentiability Properties of Functions*, Princeton University Press, Princeton, New Jersey, 1970.

[13] E. M. Stein, *Harmonic Analysis: Real Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton University Press, Princeton, New Jersey, 1993.